

Regimen Reference Order – SARC – methotrexate + vinBLAStine

ARIA: SARC – [methotrexate + vinBLAStine]

Planned Course: Once weekly until disease progression or unacceptable toxicity (28-day cycle)

Indication for Use: Advanced Aggressive Fibromatosis (Desmoid Tumor)

CVAD: Preferred (VESICANT INVOLVED)

Proceed with treatment if:

ANC equal to or greater than $1.5 \times 10^9/L$ AND Platelets equal to or greater than $100 \times 10^9/L$

❖ Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – SARC – methotrexate + vinBLAStine

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
Days 1, 8, 15 and 22		
methotrexate	30 mg/m ²	IV in normal saline 50 mL over 20 minutes (Maximum rate 10 mg/minute)
vinBLAStine	6 mg/m ²	IV in normal saline 25 mL over 5 to 10 minutes by gravity infusion

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

Day 1

- CBC, serum creatinine, urea, electrolytes and liver enzymes as per Physician Orders

Days 8, 15 and 22

- CBC

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Instruct patient to continue taking anti-emetic(s) at home
 - Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy
-

ADDITIONAL INFORMATION

- Not applicable